
RESEARCH ARTICLE

Genotypic and phenotypic correlation and path coefficient analysis for yield and other traits of sorghum (*Sorghum bicolor* L. Moench) land races at humid lowland and intermediate agro-ecology of Ethiopia

H. A. Keba, H. D. Tamir

Ethiopian Institute of Agricultural Research (EIAR), Assosa Agricultural Research Center, Postal address: 265, Assosa, Ethiopia

Corresponding authors email Id: halemu2017@gmail.com

Manuscript received: June 5, 2023; Decision on manuscript, July 2, 2023; Accepted: July 5, 2023

Abstract

This experiment was conducted to study the association among the yield contributing traits and their direct and indirect effects on the yield of sorghum. Knowing the association and path analysis of yield related traits is essential for breeders. So, this activity was conducted with the objectives of investigating the relationship among yield, yield related traits and their effect on yield. A total of 42 late and medium maturing sorghum genotypes were evaluated by using 7x6 triple lattice designs with three replications at Assosa in 2020 cropping season. For most of the investigated traits, the magnitudes of genotypic correlation coefficients were higher than the corresponding phenotypic correlation coefficients. This implies that there was inherent relationship between these traits. Grain yield showed significant and positive phenotypic and genotypic correlation with days to flowering, days to maturity, thousand grain weight and plant height. The strongest phenotypic association of grain yield was observed with thousand grain weight followed by days to maturity and days to flowering. Apart from yield greater association was recorded between days to maturity and thousand grain weight followed by disease score with insect score. The strongest positive genotypic association was observed between

grain yield and thousand grain weight followed by days to flowering and days to maturity and thousand grain weight and days to maturity. Thousand grain weights has the highest positive direct effect on grain yield while overall plant aspect has higher negative direct effect on grain yield.

Keywords: Association, direct effect, grain weight, land races, sorghum

Introduction

Sorghum (*Sorghum bicolor* (L.) Moench) crop is inherently fit to warm and arid agro-ecologies which are characterized by recurrent water deficiency and thereby challenging for growing other crops. Sorghum is the fifth most important cereal crop in the world after maize, rice, wheat and barley and only second to maize in Ethiopia. As the climatic change is becoming the major force endangering agricultural production in arid and semi-arid regions of the globe, sorghum is the front runner crop to withstand the hardship and feed the world population depends on the produce. Sorghum producers are benefitting from the crop in many ways as food, feed for livestock, porridge, cultural (local) drinks as Tella and arake which were mainly known in Ethiopia. The stalk of the crop can also serve for fencing and other buildings (Alemu and Demelash 2022).

In plant breeding, the presence of variability of the important trait which is under consideration is very important for the improvement of a crop. The primary task for plant breeders is maximizing selection efficiency to assist the identification of best genotypes with desired traits. The relationship between characters of the crop to be improved is of major importance for plant breeders. When selection is based on several plant characters, correlation analysis measures the intensity and direction of associations among characters that are important in a breeding program (Warkad *et al.*, 2010).

Studying the relationship between quantitative traits is very important for measuring the possibility of selecting two or more traits together and thereby evaluating the effect of selection for secondary traits on genetic gain for the primary trait under consideration. If the genetic correlation for both of the traits under consideration is positive, then the plant breeder can easily improve both traits at a time. It is so hard to decide the traits that fully contribute to yield when many characters are involved in trait association study because of the presence of some amount of interdependence. In this case, Path analysis is the good way of identifying main contributing character to the yield. The path coefficient analysis helps to determine the direct contribution of character and their indirect contributions via other characters (Tesfaye *et al.*, 2014). Path coefficient analysis measures the direct influence of one variable upon the other and permits separation of correlation coefficients into component of direct and indirect components. Hence, the present study is aimed to analyze and determine the traits having greater association with yield utilizing the correlation and path analysis for different traits in sorghum.

Material and methods

Forty two late and medium maturing sorghum land races collected from Wet lowland and

intermediate agro-ecologies were used for the experiment (Table1). The experiment was carried out at Assosa Agricultural Research center (AsARC) during the main season of 2019/2020 cropping season. The testing site was among the potential areas for sorghum production in the wet lowland and intermediate agro-ecological zones of the country.

Data collection and analysis

All agronomic data were collected on plot and plant bases using sorghum descriptors (IBPGR/ICRISAT, 1993). In each plot, randomly selected five plants were used to be measured for the plant based characters. The data was analyzed for Phenotypic and genotypic correlation coefficients. The observable correlation between two variables (Phenotypic correlation), which includes both genotypic and environmental components between the variables, was estimated using the formula suggested by Miller *et al.*, (1958). Path coefficient analysis involves a method of partitioning correlation coefficients into direct effect and indirect effects through alternative pathways [Pathway (P) x correlation coefficient (r)] (Aryeetey and Laing, 1973). The path coefficients were obtained by solution of simultaneous equation through the method of least square as shown by Dewey and Lu (1959). Path coefficient is calculated by solving simultaneous equations which express the basic relationship between path coefficient and correlation coefficient.

Results and discussion

Phenotypic and genotypic correlation coefficients were estimated for all possible pair of traits investigated in this trial. The results of assessment of the pair-wise associations among different agronomic traits indicated that some of the traits are positively correlated while others are negatively correlated which implies improving one traits had either positive or negative influence on the other traits.

Table1: The experimental materials used for the trial

Genotype	Genotype	Genotype	Genotype
1	ETSCAs 10001-1-1-1	25	AScol19-Kok001
2	ETSCAs 10001-1-1-2	26	Mok 079/1
3	ETSCAs 10001-1-4-1	27	AScol19-As-6
4	ETSCAs 10002-2-13-1	28	AScol19-KA021/1
5	ETSCAs 10003-3-32-1	29	Ya 036/1
6	ETSCAs 10007-2-61-1	30	Mok 079/2
7	ETSCAs 10015-2-102-1	31	AScol19-JW127
8	ETSCAs 10015-2-103-1	32	AScol19-BS 082/1
9	ETSCAs 10016-1-106-1	33	AScol19-Krm 124
10	ETSCAs 10016-1-106-2	34	AScol19-As -14
11	ETSCAs 10019-1-110-1	35	AScol19-AB126
12	ETSCAs 10019-1-115-1	36	AScol19-As-1
13	ETSCAs 10020-2-116-1	37	AScol19-SG 001
14	ETSCAs 10020-2-116-2	38	AScol19-As-13
15	ETSCAs 10020-2-116-3	39	AScol19-SG 002
16	Y039-1	40	AScol19-As-5
17	AScol19-JW128	41	Assosa-1
18	AScol19-Krm123	42	Bonsa
19	Assosa-1/1		
20	AScol19-As-7		
21	AScol19-Al25		
22	AScol19-As-2		
23	AScol19-As-8		
24	AScol19-Krm122		

Grain yield has showed highly significant ($p < 0.01$) positive phenotypic correlation with days to flowering ($r = 0.53$), days to maturity ($r = 0.63$), thousand grain weight ($r = 0.82$) and plant height ($r = 0.49$) (Table2). This implies that selection for these agronomic traits can enable to exploit higher grain yielding potential in sorghum. In the test environment, selecting sorghum genotypes with longer days to flowering and maturity, taller plant height and higher seed weight can significantly results in higher grain yield. Similar results have been reported by Alemayehu (2003) for strong positive correlation of grain yield with plant height, Mamo and Mulgeta (2020) for strong positive correlation of grain yield with thousand Kernel weight. On the other hand,

grain yield has showed highly significant ($p < 0.01$) negative correlations with over all plant aspects (PAS, $r = -0.74$), insect score, ($r = -0.33$) and bird damage ($r = -0.63$). Grain yield has showed non-significant negative association with disease score. . Strong positive phenotypic association was observed for days to flowering and days to maturity ($r = 0.71^{**}$), days to maturity and thousand grain weight ($r = 0.65^{**}$), over all plant aspects and bird damage ($r = 0.64$) while higher negative correlation was recorded for traits thousand grain weight and overall plant aspect ($r = -0.73^{**}$), thousand grain yield and bird damage ($r = -0.68$) and days to maturity and insect score ($r = -0.48^{**}$).

At genotypic level grain yield showed positive and highly significant correlations with days to flowering ($r=0.57^{**}$), days to maturity ($r=0.7^{**}$), thousand grain weight ($r=0.87^{**}$) and plant height ($r=0.53^{**}$) (Table 2). The strongest positive genotypic association was observed between grain yield and thousand grain weight ($r=0.87^{**}$) followed by days to flowering and days to maturity ($r=0.78^{**}$) and thousand grain weight and days to maturity ($r=0.78$). This showed that the positively associated characters can be improved

simultaneously and, improving one character can indirectly improve the other traits. Generally, the values of genotypic correlation coefficients were higher than the corresponding phenotypic correlation coefficients for most of the trait (Table2). This implies that there are inherent genotypic relationships between the tested traits. This result is in agreement with the previous findings of Tafere *et. al.*, (2018), Ezeaku and Mohammed (2006).

Table2: Estimates of genotypic (above diagonal) and phenotypic (below diagonal) correlation coefficients for 9 sorghum traits

	Days to flowering	Insect score	Disease score	Bird damage	Days to maturity	Grain yield	Overall plant aspect	1000 grain weight	Plant height
Insect score	1	-0.49**	-0.3ns	-0.59**	0.78**	0.57**	-0.53**	0.67**	0.54**
Insect score	-0.29**	1	0.72**	0.33*	-0.55**	-0.44**	0.4**	-0.48**	-0.42**
Disease score	-0.2ns	0.59**	1	0.16ns	-0.37*	-0.25ns	0.14ns	-0.21ns	-0.49**
Bird damage	-0.44**	0.07ns	0.02ns	1	-0.74**	-0.79**	0.76**	-0.81**	-0.52**
Days to maturity	0.71**	-0.48**	-0.3**	-0.48**	1	0.7**	-0.72**	0.78**	0.57**
Grain yield	0.53**	-0.33**	-0.19ns	-0.63**	0.63**	1	-0.78**	0.87**	0.53**
Overall plant aspect	-0.49**	0.25*	0.1ns	0.64**	-0.6**	-0.74**	1	-0.78**	-0.43**
1000 grain weight	0.6**	-0.33**	-0.16ns	-0.68**	0.65**	0.82**	-0.73**	1	0.58**
Plant height	0.54**	-0.26*	-0.37**	-0.37**	0.51**	0.49**	-0.39**	0.53**	1

*, ** significant at 5% and 1% respectively, ns= non-significant

Path coefficient analysis

Grain yield, being a dependent variable governed by many genes, is affected by many

traits. So, the phenotypic and genotypic correlations were partitioned into direct and indirect effects for the traits that are believed to have a direct relationship with grain yield from the result of this study.

Genotypic direct and indirect effects of various traits on grain yield

The estimates of genotypic direct and indirect effects of the selected traits on grain yield were presented in (Table 3). Accordingly, genotypic path analysis showed that thousand grain weights (0.57) exerted the highest positive direct effect to grain yield followed by plant height (0.04). Mengesha *et. al.*, (2019), and Chittapur and Biradar (2015) reported Similar result of direct positive correlation of plant height and thousand kernel weight on grain yield. Thousand grain weights had positive direct effect and the genotypic correlation with grain yield was positive and significant. Most of the traits scored negative direct effect on grain yield (Table3). The highest negative direct effect on grain yield was recorded for overall plant aspect or PAS (-0.21) followed by bird damage (-0.20), days to maturity (-0.07), insect score (-0.05) and days to flowering (-0.04). This implies that breeding for early maturing variety has a significant yield reduction due to the long growing period of the area. The negative direct effect of these traits has to be seriously considered because

these traits contributed to grain yield reduction in the study area. Thousand grain weights has also the highest positive indirect effect on grain yield via days to maturity (0.45), days to flowering (0.38) and plant height (0.33). These positive indirect effects contribute to increase grain yield via increasing those traits. This result is similar to the findings reported previously by Khan *et. al.*, (2013). The genotypic residual value (0.20) indicated that the traits used in the genetic path analysis explained (80%) of the variation for grain yield. In general, the trait association between yield and yield related traits in this particular study showed various magnitude of association which can be carefully considered for exploiting and selection to improve traits of interest in Ethiopian sorghum genotypes mainly land races collected from intermediate and humid lowland areas of the country. Regarding to the trait association, Singh and Chaudhary (1977) stated that whenever a trait has positive correlation and high positive indirect effects but negative direct effect on the economic trait like grain yield, emphasis should be given to the indirect effects.

Table3: Estimates of direct (bold diagonal) and indirect effect (off diagonal) at genotypic level of nine traits on grain yield

	Days to flowering	Days to maturity	Plant height	Insect score	Bird damage	Overall plant aspect	1000 seed weight	Residual
Days to flowering	-0.04	-0.06	0.02	0.03	0.12	0.11	0.38	0.20
Days to maturity	-0.03	-0.07	0.02	0.03	0.15	0.15	0.45	0.20
Plant height	-0.02	-0.04	0.04	0.02	0.10	0.09	0.33	0.20
Insect score	0.02	0.04	-0.02	-0.05	-0.07	-0.09	-0.28	0.20
Bird damage	0.02	0.05	-0.02	-0.02	-0.20	-0.16	-0.46	0.20
Overall plant aspect	0.02	0.05	-0.02	-0.02	-0.15	-0.21	-0.45	0.20
1000 seed weight	-0.02	-0.06	0.02	0.03	0.16	0.17	0.57	0.20

R-square =0.80

In conclusion, knowledge of the relationships among different agronomic traits as well as their association with grain yield is of paramount importance in crop improvement. Grain yield is found to be positively and significantly associated with days to flowering, days to maturity, thousand grain yields and plant height at both phenotypic and genotypic level. So, positive genotypic correlation implies that the traits are associated inherently.

References

1. Tesfaye ,W.M., Adugna, W. and Tsige ,G. 2014. Correlation and path coefficient analysis among yield component traits of Ethiopian mustard (*Brassica Carinata* a. Brun) at Adet, Northwestern. Ethiopia. J. Plant Sci., 2(2): 89-96
2. IBPGR/ICRISAT 1993. Descriptors for sorghum. IBPGR Secretariat, FAO, Rome, Italy. pp.1-26.
3. Miller, P.A., Williams, J. C., Robinson, H.F., and Comstock, R.E. 1958. Estimate of genotypic and environmental variances and co-variance in upland cotton and the implication selection. Agrono. J., 50 (3): 126-131.
4. Areyetey, A.N. and Laing, E. 1973. Inheritance of yield components and their correlation with yield in cowpea. (*Vigna unguiculata* (L) Walp.). Euphytica, 22: 386-392.
5. Dewey, R.R. and Lu, K.H. 1959. A correlation coefficient analysis of components of crested seed production. Agrono. J., 51 (9): 515- 518.
6. Alemayehu, A. (2003). Genetic variability and breeding potential of barley (*Hordeum vulgare* L.) landraces from North Shewa in Ethiopia,” PhD Thesis, Faculty of Natural and Agricultural Sciences University of Free State, Bloemfontein, South Africa. 226.
7. Mamo, M. and Worede, F. 2020. Character association and Path analysis of sorghum (*Sorghum bicolor* (L.) Moench) genotypes for Striga resistance. B S J Agri., 3(2): 96-102
8. Ezeaku, I.E. and Mohammed, S.G.2006. Character association and path analysis in grain sorghum. African J. Biotech., 5(14):1337- 1340.
9. Tafere, M., Sentayehu ,A., Taye, T. ,and Dagne, W. 2018. Correlation and path coefficient analysis for agronomical traits of lowland adapted Ethiopian sorghum genotypes [*Sorghum bicolor* (L.) Moench] Genotypes. J. Bio., Agril. Healthcare, 8(15):29-32.
10. Chittapur, R. and Biradar, B.D. 2015. Association studies between quantitative and qualitative traits in rabi sorghum. Indian J. Agric. Sci., 49(5): 468–471.
11. Mengesha ,G.H., Hailemariam, F.M., Mindaye T.T., Lakew, B. and Verma, R.P.S. 2019. Correlation and path analysis of yield, yield contributing and malt quality traits of Ethiopian sorghum (*Sorghum bicolor* (L.) Moench) genotypes. African J. Plant Sci., 13(8): 209-220,
12. Singh, R.K. and Chaudhary, B.D.1977. Biometrical Methods in Quantitative Genetic Analysis. Kalyani Publishers, New Delhi, pp.318 .
13. Khan, H.A., Shad, S.A. and Akram, W. 2013. Resistance to new chemical insecticides in the house fly, *Musca domestica* L., from dairies in Punjab, Pakistan. Parasitology Res., 112 (5):2049-54.
14. Warkad, Y.N., Tidke, R.T., Maske, N.M., Kharde, A.V. ,and Potdukhe, N.R.2010. Character association and path analysis in sorghum [*Sorghum bicolor* (L.)] Moench. Internat. J. Agric. Sci., 6 (1):100-104.