
RESEARCH ARTICLE

Studies on genetic variability and correlation for seedling vigour traits in rice (*Oryza sativa* L.) genotypes

M. Rathia¹, S. Kar¹, R.R. Kanwar¹, N.C. Mandavi¹, G.Thakur¹, V.Dansena²

1 S.G. College of Agriculture and Research Station Kumharawand, Jagdalpur, Bastar 494 001 Chhattisgarh, India

2 Krantikari Debridhur College of Horticulture and Research Station, Jagdalpur, Bastar 494001 Chhattisgarh, India

Corresponding authors email: minakshirathia2905@gmail.com

Manuscript received: December 1, 2024; Decision on manuscript, December 30, 2024; Manuscript accepted: January 1, 2025

Abstract

Seedling vigour is important agronomical traits and contributor to subsequent tillering quality and yield of rice. Crop stand is dependent on the seedling vigour. For higher grain yield, good crop stand is equally important along with high yield potential. Therefore the present study was carried out to explore the genetic variability and correlation among eleven seedling vigour related traits in thirty-three rice genotypes including three checks viz., Narendra 359, Karma mahsuri and IGKVR 1244, at S.G. College of Agriculture and Research Station, Jagdalpur, Chhattisgarh, India. High heritability was found for all the eleven seedling vigour traits. Highest seedling vigour-I at 14th day was recorded for Narendra 359 followed by R2699-31-1, R2326-108-1-61-1 and R2737-25-1. High heritability coupled with high GAM was recorded for root/shoot ratio at 7th day, shoot length at 14th day, shoot length at 7th day and root/shoot ratio at 14th day. The findings from the character association analysis revealed that seedling vigour index-I at 14th day showed high positive significant association with characters like germination percentage, shoot length at 14th day, root length at 14th day and seedling height at 14th day. Further, these promising

genotypes can be used in future breeding programme.

Keywords: Rice, seedling vigour, variability, heritability, correlation

Introduction

Rice is the most important cereal of the world providing 21% of global human per capita energy and 15% of per capita protein. Rice (genus *Oryza*) has only two domesticated species *Oryza sativa* L. and *Oryza glaberrima* L., out of its twenty two species. Hence, *Oryza sativa* L. is the major food crop for people in Asia and nearly 90% of the world's rice is produced and consumed in this region (Warusawithana *et al.*, 2017). In world total rice grown in area 165.68 million hectares with the production of 514.57 million tonnes and productivity of 4640 kg per hectare during 2022-23 (Anonymous, 2024). In the world, 85% of the total rice area is in Asia. Asia is the world's leading producer of rice, accounting for around 90% of global output. People in Asian countries consume more than 75% of the world's rice, making rice vital to Asia's food security. Because of the predicted growth in the population, rice demand is expected to rise even more (Tripathi *et al.*, 2011).

India is the second largest producer and consumer of rice in the world. Rice breeding in India has made significant impact in the socioeconomic spheres by simultaneously ensuring food security as well as commerce. In India total rice grown in area 47.83 million hectares with the production of rice during 2022-23 is estimated at record 135.75 million tonnes and productivity of 2838 kg per hectares (Anonymous, 2023). Chhattisgarh is known as “rice bowl of India” and about 82% population of the state is dependent on agriculture for their livelihood. In Chhattisgarh total rice grown area is 3.77 million hectares with the production of 9.81 million tonnes and productivity of 2602 kg per hectare during 2022-23 (Anonymous, 2023). Rice seedling vigour is a complex quantitative feature that has been encoded in nature. Seedling vigour is an agronomical traits and sign of potential seed germination. It improves the speed, uniformity of the seedlings with stand establishment and grain yield. It determines the quick, uniform emergence and growth of seedlings (Anumalla *et al.*, 2015). Seedling vigour is an important characteristic of seed quality, reflecting potential seed germination, seedling growth, seed longevity and tolerance to adversity. Seedling with strong vigour may significantly improve the speed and uniformity of seed germination and lead to perfect field emergence, good crop performance, and even high yield under different conditions (Sun *et al.*, 2007). For higher seed yield, good crop stand is equally important along with high yield potential. Crop stand is dependent on the seedling vigour. Some characters like root length, shoot length and seedling height has been identified as standard indicator of seedling vigour and seedling vigour related a trait which belongs to *Oryza sativa* L. ssp. indica (Akshaya *et al.*, 2020). Hence, the current study was initiated with the main aim of identifying rice genotypes with superior early seedling vigour related traits which belongs to *Oryza sativa* L. ssp. Indica.

Materials and method

The experiment was conducted during *Kharif* 2023. Thirty three genotypes of rice were evaluated for seedling vigour traits in randomized block design with two replications at Research cum Instructional Farm of S.G. College of Agriculture and Research Station, Kumhrawand located in Jagdalpur, Bastar (Chhattisgarh, India). All laboratory observations were recorded in each replication was considered for statistical analysis. Estimation of seedling vigour traits In order to assess and quantify the genetic variability among the genotypes for the characters under study, the following parameters were estimated as given below. Germination percentage (%), root length (cm) at 7th day, shoot length (cm) at 7th day, seedling height (cm) at 7th day, root/shoot ratio at 7th day, seedling vigour index-I at 7th day, root length (cm) at 14th day, shoot length (cm) at 14th day, seedling height (cm) at 14th day, root/shoot ratio at 14th day, seedling vigour index-I at 14th day recorded as suggested by Sai Teja *et al.*, (2024).

Results and discussion

In ANOVA all the characters were found to be significant at 1% level (Table not shown). The presence of highly significant values of mean sum of square for all the 11 characters indicated that high variability was present among the genotypes for the characters. Table 1 presents the results of the genetic variability parameters, including mean, genotypic coefficient of variability (GCV), phenotypic coefficient of variability (PCV), broad sense heritability (h^2), and genetic advance as a percentage of mean (GAM), for seedling vigour characteristics. The magnitude of PCV values for all traits was greater than the corresponding GCV values, suggesting that these characters may be influenced by the environment. Phenotypic coefficient of variation values was found to be slightly higher than genotypic coefficient of variation values showing the influence of environment on trait expression.

Table 1: Genetic variability parameters of 11 studied characters

Traits	Mean	Minimum	Maximum	Heritability (%)	GA mean (%)	GCV (%)	PCV (%)
Germination (%)	94.3	89.1	99.0	76.2	3.7	2.1	2.4
Shoot length 7 th day	2.3	1.8	3.2	98.2	24.7	12.1	12.2
Root length 7 th day	9.1	6.6	10.9	91.7	16.8	8.5	8.9
Root /shoot 7 th day	4.0	3.0	5.6	95.8	25.6	12.7	13.0
Seedling height 7 th day	11.3	9.00	13.3	66.8	13.6	8.0	9.8
Seedling vigour index-I 7 th day	10.7	8.6	12.4	68.7	14.4	8.4	10.2
Shoot length 14 th day	7.1	5.3	9.2	94.6	25.0	12.5	12.8
Root length 14 th day	17.4	14.4	21.8	91.5	14.0	7.1	7.4
Root /shoot 14 th day	2.53	1.91	3.1	86.9	24.0	12.5	13.4
Seedling height 14 th day	24.5	20.1	29.5	95.1	14.4	7.1	7.3
Seedling vigour index-I 14 th day	23.2	18.7	28.8	96.0	16.1	7.9	8.1

Table 2: Phenotypic correlation coefficients of yield and yield attributing characters

Traits	Germi nation (%)	Shoot length 7 th day	Root length 7 th day	Root /shoot 7 th day	Seedlin g height 7 th day	Seedling vigour index-I 7 th day	Shoot lengt h 14 th day	Root lengt h 14 th day	Root /shoot 14 th day	Seedlin g height 14 th day	Seedlin g vigour index-I 14 th day
Germination (%)	1.000	0.128	-0.07	-0.166	0.02	0.23	0.16	0.15	-0.08	0.19	0.43**
Shoot length 7 th day			0.275*	-0.717**	0.52**	0.54**	-0.09	-0.21	-0.03	-0.20	-0.15
Root length 7 th day				0.444**	0.82**	0.78**	-0.26*	-0.19	0.14	-0.27*	-0.27*
Root /shoot 7 th day					0.08	0.04	0.00	0.08	0.05	0.06	0.00
Seedling height 7 th day						0.97**	-0.20	-0.23	0.07	-0.27*	-0.24*
Seedling vigour index-I 7 th day							-0.15	-0.19	0.04	-0.22	-0.13
Shoot length 14 th day								0.31*	-0.82**	0.73**	0.71**
Root length 14 th day									0.25*	0.87**	0.84**
Root /shoot ratio 14 th day										-0.23	-0.23
Seedling height 14 th day											0.96**
Seedling vigour index-I 14 th day											1.00

*, ** significant at 5% and 1% level, respectively

Fig. 1: Bar diagram showing heritability coupled with genetic advance as percent of mean for yield and other traits




Fig.2: Bar diagram showing phenotypic and genotypic coefficient of variation for yield and other traits




Fig. 1 represents the bar diagram showing heritability coupled with genetic advance as percent of mean for yield and yield attributing traits. Moderate GCV and PCV values were observed for characters like shoot length at 7th day, root/shoot ratio at 7th day, shoot length at 14th day and root/shoot ratio at 14th day. High heritability was found for all the 11 seedling vigour traits and supported by earlier findings of Singh *et al.*, (2024). High heritability coupled with high GAM was recorded for root/shoot ratio at 7th day, shoot length at 14th day, shoot length at 7th day and root/shoot ratio at 14th day. Similar results were obtained for germination percentage, seedling height, seedling vigour index-I, root length and shoot length by Mounika Reddy *et al.*, (2022). Bezawada *et al.*, (2020) reported similar results for germination percentage, seedling height, shoots length and root length. The findings from the character correlation analysis revealed that seedling vigour index-I at 14th day showed high positive significant association with characters like germination percentage, shoot length at 14th day, root length at 14th day and seedling height at 14th day (Table 2). Fig.2 represents the bar diagram showing phenotypic and genotypic coefficient of variation for yield and yield attributing characters. Further, these promising genotypes

can be used in future breeding programme. Similar results were obtained by Teja *et al.*, (2024) for traits like shoot length at 7th day, root length at 7th and 14th day.

Hence, in conclusion genotype Narendra359 followed by R2699-31-1, R2326-108-1-61-1 and R2737-25-1 recorded good seedling vigour index. Moderate GCV and PCV values were reported for characters like shoot length at 7th day, root/shoot ratio at 7th day, shoot length at 14th day and root/shoot ratio at 14th day. High heritability was found for all the 11 seedling vigour traits. Highest heritability combined with high GAM was recorded for root/shoot ratio at 7th day, shoot length at 14th day, shoot length at 7th day and root/shoot ratio at 14th day. The findings of this study can be used for future breeding programs to develop varieties that are not only high-yielding but also have strong seedling vigour, which is important for successful crop establishment and better overall crop performance.

Acknowledgements

The corresponding author is thankful to SPVT and S. G. College of Agriculture and Research Station, Kumharawand, Jagdalpur, IGKV Raipur, Chhattisgarh for providing facility to work under this research project.

References

1. Akshaya, M., Thirumurugan, T., Chitra, S., Nithila, S., and Jeyaprakash, P. 2020. Genetic variability in rice (*Oryza sativa* L.) landraces for seedling vigour traits. *Electronic J. Plant Breed.*, 11(1): 91-96.
2. Anonymous, 2023. UPAg, Unified Portal for Agricultural Statistics, Department of Agriculture and Farmers Welfare, Govt. of India, New Delhi.
3. Anonymous, 2024. World Agricultural Production, Foreign Agricultural Service, United States Department of Agriculture, Washington, D.C., United States, March, p. 28.
4. Anumalla, M., Anandan, A. and Sharat, P. K. 2015. Early seedling vigour, an imperative trait for direct-seeded rice: an overview on physio- morphological parameters and molecular markers. *Planta*, 241:1027–1050.
5. Bezawada, A., Senguttuvei, P., Hema Latha, V., Yamini, K.N., Jhansi rani, K. and Beulah, P. 2020. Variability and correlation analysis for seedling vigour traits in rice (*Oryza sativa* L.) genotypes. *Int.J. Current Microb. Applied Sci.*, 9(7):2877-2887.

6. Singh, D.P. Indrajeet, Kar, S., Kanwar, R.R. 2024. Large scale evaluation, genetic variability and correlation analysis in rice accessions. J. Genet. Genom. Plant Breed., 8 (3): 83-91.
7. Reddy, Y.M., Vasanthi, R.P., Reddy, B.R.K., Murthy, B.R. and Vemireddy, L.R. 2022. Genetic Parameters for quantitative and early seedling vigour related traits in rice (*Oryza Sativa* L.) under dry direct seeded rice conditions. Andhra Pradesh J. Agril. Sci., 8(3): 170-173.
8. Teja, S. D., Balakrishanan, D., Kar, S., Salam, J.L., Singh, D.P., Kumar, A.P., Swamy, A.V.S.R. and Subba Rao, L.V. 2023. Genetic variability and correlation studies in high yielding varieties of rice (*Oryza Sativa* L.). J. Rice Res.. 16 (2). 1-13.
9. Sun, Q., Wang, J.H. and Sun, B.Q. 2007. Advances on seed vigor physiological and genetic mechanisms. Agric. Sci. China.6 (9):1060–1066.
10. Tripathi, K.K., Warriar, R., Govila, O.P. and Ahuja, V. 2011. Biology of *Oryza sativa* L.(rice). India: Series of Crop Specific Biology Documents. Ministry of Science and Technology, India.
11. Warusawithana, T. M., Samarasinghe, W. L. G., Dasanayaka, P. N., Ubeysekara, N. M. and Jayarathna, K. G. C. N. 2017. Genetic diversity analysis of traditional Rice variety Pachchaperumal using SSR markers. Int. J. Sci. Tech. Res., 6(8): 229-237.